

2



AD-A238 084



THE SOLD

DTIC
JUL 15 1991

INSTITUTE REPORT NO. 461

COMPARISON OF INTRAOSSEOUS AND INTRAVENOUS
DELIVERY OF HYPERTONIC SALINE/DEXTRAN (HSD) IN
ANESTHETIZED, EUVOLEMIC PIGS


M.A. Dubick
J.W. Pfeiffer,
C.B. Clifford,
D.E. Runyon, and
G.C. Kramer

PRELIMINARY STATEMENT A

Approved for public release;
Distribution Unlimited

Division of Military Trauma Research

April 1991



LETTERMAN ARMY INSTITUTE OF RESEARCH PRESIDIO OF SAN FRANCISCO CALIFORNIA 94129

33

91-04904



91 7 12 033

**Comparison of Introsseous and Intravenous Delivery of Hypertonic Saline/Dextran (HSD)
in Anesthetized, Euvoletic Pigs, M.A. Dubick et al.**


This document has been approved for public release and sale; its distribution is unlimited.

Destroy this report when it is no longer needed. Do not return to the originator.

Citation of trade names in this report does not constitute an official endorsement or approval of the use of such items.

The experimental studies of the author described in this report were reviewed and approved by the Institutional Review Committee/Animal Care and Use Committee at Letterman Army Institute of Research. The Manuscript was peer reviewed for compliance prior to submission for publication. In conducting the research described here, the author adhered to the "Guide for the Care and Use of Laboratory Animals," DHEW Publication (NIH) 85-23.

This material has been reviewed by Letterman Army Institute of Research and there is no objection to its presentation and/or publication. The opinions or assertions contained herein are the private views of the author(s) and are not to be construed as official or as reflecting the views of the Department of the Army or the Department of Defense. (AR 360-5)

 17 April 91
DONALD G. CORBY (date)
COL, MC
Commander

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
1a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release Distribution is unlimited.		
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE					
4. PERFORMING ORGANIZATION REPORT NUMBER(S) Inst Report No. 461			5. MONITORING ORGANIZATION REPORT NUMBER(S)		
6a. NAME OF PERFORMING ORGANIZATION Letterman Army Institute of Research		6b. OFFICE SYMBOL (If applicable) SGRD-ULT-M	7a. NAME OF MONITORING ORGANIZATION US Army Medical Research and Development Command		
6c. ADDRESS (City, State, and ZIP Code) Division of Military Trauma Research LAIR Presidio of San Francisco, CA 94219-6800			7b. ADDRESS (City, State, and ZIP Code) Ft. Detrick, Frederick, MD 21701		
8a. NAME OF FUNDING/SPONSORING ORGANIZATION		8b. OFFICE SYMBOL (If applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER		
8c. ADDRESS (City, State, and ZIP Code)			10. SOURCE OF FUNDING NUMBERS		
			PROGRAM ELEMENT NO. 63807A	PROJECT NO. D836	TASK NO. AX
11. TITLE (Include Security Classification) Comparison of Intraosseous and Intravenous Delivery of Hypertonic Saline/Dextran (HSD) in Anesthetized, Euvoletic Pigs					
12. PERSONAL AUTHOR(S) M.A. Dubick, J.W. Pfeiffer, C.B. Clifford, D.E. Runyon, G.C. Kramer					
13a. TYPE OF REPORT Institute		13b. TIME COVERED FROM _____ TO _____	14. DATE OF REPORT (Year, Month, Day) 1991 April 1		15. PAGE COUNT 17
16. SUPPLEMENTARY NOTATION					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number) (U) intraosseous, hypertonic resuscitation, dextran, pigs (U)		
FIELD	GROUP	SUB-GROUP			
19. ABSTRACT (Continue on reverse if necessary and identify by block number) Delivery of fluids and drugs by intraosseous (IO) infusion has seen renewed use in pediatrics and has been proposed for adult use. To evaluate the efficacy of IO infusion of resuscitation fluids in adults, the present study investigates vascular entry cardiodynamic (CD) responses to IO and IV infusion of the small volume resuscitation formulation, 7.5% NaCl/6% Dextran-70 (HSD) in 12 anesthetized, euvoletic pigs. Using a sternal IO access device, 4 ml/kg HSD was infused over 2-6 min; each IO infusion was paired with an IV infusion of equal time. IO and IV HSD infusion produced similar CD effects and increased plasma volume over 20% for the 2 h experimental period. Delivery of Na and dextran was essentially complete within 1 min after infusion and plasma Na and dextran concentrations were comparable over the 2 h period. Histological examination of marrow showed a small, focal hemorrhage as the predominant effect of IO infusion of HSD.					
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT <input type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED		
22a. NAME OF RESPONSIBLE INDIVIDUAL Donald G. Corby, COL, MC			22b. TELEPHONE (Include Area Code) (415) 561-3600		22c. OFFICE SYMBOL SGRD-ULZ

These data indicate that IO delivery of HSD is as rapid as IV delivery, and may be a viable alternative for emergency resuscitation where vascular access is compromised.

**Comparison of Intraosseous and
Intravenous Delivery of Hypertonic
Saline/Dextran (HSD) in Anesthetized,
Euvolemic Pigs**

Michael A. Dubick, Ph.D., Juergen W. Pfeiffer, M.S.,
Charles B. Clifford, D.V.M., Ph.D., David E. Runyon, B.S.,
and George C. Kramer, Ph.D.¹

Division of Military Trauma Research
Letterman Army Institute of Research
Presidio of San Francisco, CA 94129-6800

¹ Division of Anesthesia Research
University of Texas Medical Branch
Galveston, TX 77550

Accession For	
DTIC 02421	
DTIC 122	
Unannounced	
Justification	
6.	
Distribution	
Availability: Codon	
Dist	Avail and/or Special
A-1	

ABSTRACT

Delivery of fluids and drugs by intraosseous (IO) infusion has seen renewed use in pediatrics and has been proposed for adult use. To evaluate the efficacy of IO infusion of resuscitation fluids in adults, the present study investigates vascular entry cardiodynamic (CD) responses to IO and IV infusion of the small volume resuscitation formulation, 7.5% NaCl/6% Dextran-70 (HSD) in 12 anesthetized, euvolemic pigs. Using a sternal IO access device, 4 ml/kg HSD was infused over 2-6 min; each IO infusion was paired with an IV infusion of equal time. IO and IV HSD infusion produced similar CD effects and increased plasma volume over 20% for the 2 h experimental period. Delivery of Na and dextran was essentially complete within 1 min after infusion and plasma Na and dextran concentrations were comparable over the 2 h period. Histological examination of marrow showed a small, focal hemorrhage as the predominant effect of IO infusion of HSD. These data indicate that IO delivery of HSD is as rapid as IV delivery, and may be a viable alternative for emergency resuscitation where vascular access is compromised.

Comparison of Intraosseous and Intravenous Delivery of
Hypertonic Saline/Dextran (HSD) in Anesthetized, Euvolemic
Pigs -- Dubick et al.

Introduction

In recent years, a number of reviews and reports have re-introduced the use of intraosseous (IO) infusion of fluids and drugs in emergency situations (1-7). The emphasis of IO infusion in these previous studies concerned initial resuscitation and drug delivery in childhood emergencies. It is apparent, however, that IO infusion into red bone marrow could be utilized for rapid vascular access in patients of all ages, under circumstance where securing a peripheral or central intravenous (IV) line is difficult or delayed.

The majority of studies employing IO infusions have used the tibia as the site of infusion (7-12), although the sternum has also been used (11-12). In pediatric patients the proximal tibia is the preferred site for IO access (5). However, in adults the red marrow of long bones is replaced with relatively avascular yellow or fatty marrow. Thus, in adults the sternum or bones of the pelvic girdle are the main sites of red marrow, and have been used for IO infusion (11-12).

In our ongoing investigations into the efficacy of 7.5% NaCl/6% Dextran-70 (HSD) for treatment of hypovolemia, we have begun evaluating sternal infusion of HSD as a viable alternative in field resuscitation. The sternum's large marrow space of uniform geometry is a definite advantage, along with its lying under only a thin layer of skin. Consequently, an IO access device was developed to provide rapid, reliable vascular access through the sternum (13).

In preliminary studies with this device, IO infusion of 200 ml of HSD over 2 to 4 min was used to successfully resuscitate sheep from moderate hemorrhagic shock (14,15). While IO infusion of HSD was rapidly effective in resuscitation from hemorrhagic hypotension (14,15), IO infusion of normal saline (NS) was more slowly effective, as vascular entry of the much larger volume of NS required for resuscitation was impeded by the marrow's naturally high resistance to flow. Encouraged by these preliminary studies with IO delivery of HSD, the present study evaluates the rate of vascular entry of HSD using IO infusion in euvolemic swine and compares with data following IV infusion. Additionally, lungs and sternum were evaluated to detect any significant acute histopathology associated with the IO infusion of HSD.

Materials and Methods

Animals and Treatment

Twelve immature Yorkshire swine (33.7 ± 2.7 kg) were obtained from the colony at the University of California, Davis. Animals were housed individually in an indoor laboratory holding facility with a 12 h light/dark cycle, were maintained at constant temperature and humidity, and were fed commercial chow and water ad libitum.

Animals were fasted 24 hours before surgery. Swine were premedicated with ketamine HCl (~ 2 mg/kg), xylazine HCl (~ 2 mg/kg) and atropine (~ 0.1 mg/kg) and then anesthetized with halothane (1-2%), nitrous oxide 50%, and oxygen for the entire experiment. Pigs were allowed to breathe spontaneously. Catheters were placed in the aorta and vena cava via the carotid artery and jugular vein for infusions and pressure monitoring as needed. To monitor central venous and pulmonary artery pressures and to determine cardiac output, a 7 French Swan Ganz thermal dilution catheter was placed into the pulmonary artery via the internal jugular vein. The femoral artery was catheterized for bleeding and blood sampling. A baseline period of 60 minutes was allowed before beginning each experiment. Pigs remained anesthetized throughout the entire experimental period.

After the baseline period a 4 ml/kg dose of 7.5% NaCl/6% Dextran 70 (HSD; Pharmacia AB, Uppsala, Sweden) was infused into either the manubrium ($n=6$) or the jugular vein ($n=6$). Blood samples were taken at 5, 30 and 60 minutes before infusion and at 1, 2, 4, 6, 10, 15, 30, 45, 60, and 120 minutes after the end of the infusion. The times required to infuse HSD IO were noted, and paired with an IV experiment of identical infusion time.

Physiological Measurements

A multichannel monitor continuously recorded arterial pressure, central venous pressure, pulmonary arterial pressure, heart rate and ECG throughout the experiments. Systemic arterial pressure, heart rate, central venous pressure, pulmonary artery and wedge pressure were recorded at 3 baseline time periods and at 1, 2, 4, 10, 15, 30, 60, and 120 minutes after infusion. Cardiac output was

determined by thermodilution at 2, 10, 15, 30, 60 and 120 minutes after infusion. Blood samples taken from the arterial line were used for biochemical analysis and determination of plasma volume.

Biochemical Assays

Total carbohydrate concentrations in plasma were determined by the anthrone reaction (16) following precipitation of serum protein with 10% trichloroacetic acid (TCA). Plasma glucose was determined by an automated glucose-hexokinase enzymatic method performed by the Analytical Chemistry Branch, Letterman Army Institute of Research (LAIR). Plasma dextran concentrations were then estimated by subtracting the glucose concentrations from the concentrations of total carbohydrate.

Assays for sodium, potassium, chloride and protein were also performed by the Analytical Chemistry Branch, LAIR. Plasma sodium and potassium concentrations were determined with a flame photometer (Instrumentation Laboratory, Lexington, MA). Chloride and protein concentrations were determined by a commercial kit (Roche Diagnostic Systems, Nutley, NJ) and the Biuret method, respectively, both adapted for analysis on a Cobas Fara II centrifugal fast analyzer (Roche Analytical Instruments, Belleville, NJ).

Other assays

Plasma volume was determined at baseline using the Evans blue dye dilution method (17). Plasma volume expansion after infusion of HSD was calculated from the baseline plasma volume and the decrease in plasma protein concentration or hematocrit (18).

Histological Evaluation

All animals were examined at necropsy. Tissues were saved in 10% neutral buffered formalin. Lungs were inflated at a pressure of approximately 30 cm water with formalin. A cross section of sternum at the injection site, and liver, kidney, spleen, and heart or any gross lesions were routinely examined by light microscopy. In addition, six sagittal sections of the left inferior lobe of the lung, plus lung sections from any gross pulmonary lesions were examined by light microscopy. All light microscopy was of 4-6 micron thick paraffin sections stained with hematoxylin

and eosin. Any histologic lesions observed were graded on a scale of 1 to 5, where 1= minimal, 2= mild, 3= moderate, 4= marked, and 5= severe. A particular focus of the inspection of lung tissue was evidence for thrombi, fat or bone emboli, necrosis or hemorrhage. Similarly, the sternal sections were examined for decreased cellularity (washout) or hemorrhage.

Statistical Analysis

Data were statistically analyzed by 2-way ANOVA with time and either infusion route or infusion time as the variables (19). Data were also analyzed by co-variate analysis (19) to adjust for differences in infusion time between the two groups. The ANOVA was corrected for repeated measures on the time factor. A $p < 0.05$ was taken as the level of significance.

Results

In two of the pigs, the IO infusion of 4 ml/kg HSD into the sternum required 2 min; the remaining pigs required 3, 4, 5, and 6 min each. Therefore, to insure proper matching of infusion times, 4 ml/kg HSD was administered IV to an equal number of pigs in times corresponding to the IO infusion.

Baseline values for mean arterial pressure (MAP) and cardiac output (CO) were similar for all pigs prior to HSD infusion (Figure 1). IO and IV infusion induced an immediate, transient decrease in MAP, typically observed in anesthetized pigs, that returned to pre-infusion pressure 1 to 4 min after infusion (Figure 1A). No significant differences were observed in MAP between the IO and IV groups throughout the 2 h experimental period.

In pigs infused IO with HSD, CO was increased 18% over baseline values at the end of infusion (independent of the infusion time) and remained about 20% above baseline throughout the experimental period (Figure 1B). CO appeared less augmented following IV HSD infusion, but the differences were not statistically significant between the two groups (Figure 1B). It appeared that the greater augmentation in CO in the IO group was associated with an animal infused over 4 min. If this animal was eliminated

from the analyses, IO and IV infusion of HSD would have a more similar effect on CO. This pig, however, was retained in the analyses for completeness and because it did not affect the overall data interpretation.

Plasma volume in euvolemic pigs was expanded over 20% with both IO and IV infusion of HSD, as determined by Evan's Blue dye dilution (Figure 1C). Changes in hematocrit and plasma protein concentrations (Table 1) following HSD infusion also confirmed a similar expansion of plasma volume in these animals. Plasma volume expansion by either route of HSD administration was also independent of the 2-6 min infusion time.

Both IO and IV infusion of HSD resulted in a rapid rise in plasma dextran and Na concentrations 1 to 2 min following infusion (Fig. 2, Table 2). Although Na concentrations were significantly higher ($p < 0.05$) at the end of the 2 h post-infusion period in comparison with baseline concentrations, the differences were not considered of clinical significance (Table 2). In contrast to Na, plasma dextran concentrations dropped more slowly during the post-infusion period and no significant differences were observed in dextran concentrations following either route of administration, suggesting no effect of infusion route on either dextran entry or the early phases of vascular clearance of dextran (Fig. 2). Peak Na and dextran concentrations also appeared to be independent of the varying infusion times encountered in this study, i.e., peak concentrations of Na or dextran occurred 1-2 min after the end of infusion. In addition, the method of infusion did not significantly affect plasma K^+ concentrations, although Cl^- rose immediately after infusion and then slowly returned towards pre-infusion levels (Table 2).

Total vascular dextran levels were calculated from plasma concentrations and plasma volume estimates to further evaluate delivery of dextran into the circulation. Estimated vascular dextran peaked at the end of the infusion period or at 1 min thereafter. Although estimated peak total vascular dextran tended to be higher in the IV than IO group, the differences were not statistically significant. It should be noted that the amount of dextran infused in the IO group (7.66 ± 0.50 g) was not significantly different from the amount infused in the IV group (7.14 ± 0.66 g). In both groups the data suggest that the total dose of dextran was accounted for in the vascular space and decreased about 25% after 2 h in both groups (data not shown). Thus, both

groups received similar amounts of dextran and the data obtained were again independent of infusion times.

Pathological Evaluation

As shown in Table 3, an equal number of pigs in each group had a minimal degree of acute to subacute pulmonary inflammation. Evidence for pulmonary thrombi or emboli was observed in both groups of pigs. When present, these emboli were small ($<30\ \mu$, diameter) and did not appear to occlude the vessels in which they were found, nor were they associated with lesions in the pulmonary vessels or parenchyma. In no case did the emboli consist of bone marrow components. Examination of the sternum revealed evidence of hemorrhage in 5/6 pigs infused IO with compared with 2/6 pigs infused IV (Table 3). Washout of sternal cellular marrow elements was observed in 1 pig in each group.

Discussion

The data from the present study indicate that IO infusion of HSD resulted in rapid vascular entry of Na and dextran, plasma volume expansion and augmentation of CO; i.e., all measures of sternal IO infusion were as efficacious as IV infusion. Using covariate analysis, these results were found to be independent of the 2 to 6 min infusion times required for IO administration of HSD into these pigs. Previous studies using IO infusion of various drugs have reported rapid distribution of the infused substance throughout the general circulation (12,20-22). Early studies by Tocantis, et al, (12) showed that a dye solution infused in the sternum leaves the marrow and enters the right heart very quickly. The vascular Na and dextran data from the present study are in agreement with these observations since plasma Na and dextran concentrations peaked by the end of infusion or within a minute thereafter. In addition, calculation of the estimated vascular dextran support complete delivery into the vasculature of the dextran infused IO.

In their studies of IO infusion of Ringer's lactate for the treatment of hypovolemia in dogs, Hodge, et al. (23) suggested that due to limitations in flow and fluid volumes

required, sufficient flow rates may not be attained for definitive treatment of hemorrhagic shock. Thus, the reduced volume requirements for effective resuscitation with HSD (18,24,25) make it a potential useful fluid for IO infusion. This is further supported by the observation that the infusion times in the present study are low and within the limits of infusion times employed for IV HSD administration (18,25). Recently, Chevez-Negrete, et al. (26,27) demonstrated that the adult sternum can be successfully used as a site for emergency resuscitation with 250 ml of HSD. Emergency room treatment of patients hypotensive from GI bleeding were treated more effectively with IO infusion of HSD than with IV infusion of crystalloids and blood (27). A third group treated with HSD IV, has a similar outcome to those infused IO.

The use of IO infusion methods has been limited by concerns of adverse reactions. Although a number of adverse reactions have been reported following IO infusion into both the tibia and the sternum (1,28), the complication rate of IO infusion appears no greater than that of IV infusion; the consequences of IO complications, however, can be greater. In particular, the effects of localized infection are often significantly more severe in bone than in vein and surrounding soft tissues (1,5,6). Fortunately, osteomyelitis is relatively rare after IO infusion, (1,5) suggesting that aseptic techniques are practical and effective. In addition, extravasation of infused fluids into the soft tissue over the sternum or into the pleural space are relatively rare, but reported complications (5). In the present study, we employed a new sternum needle designed to minimize both back flow along the needle shaft and to prevent puncture of the inner cortical bone (13).

As mentioned, the hydraulic resistance of red marrow is high, requiring greater infusion pressures than those generated by standard IV gravity drip or IV pressure bag (300 mm Hg) (8). In the present study, we used syringe delivery (1-2 atm pressure) to infuse 4 ml/kg HSD. Such pressurized delivery of fluids into the sternum is potentially dangerous because if the IO needle becomes dislodged or is not tightly imbedded in the bone, then soft tissue extravasation can occur. The needle used in the present study has a bulbous threaded head which lies entirely in the marrow. A wide, circular base and spring are designed to pull up on the bulbous head, seal the puncture hole, and stabilize the needle (13,15). In the

present study, sternal infusion of 4 ml/kg HSD did not cause clinically significant pulmonary bone or fat emboli, while marrow examinations showed only a small focal hemorrhage as the predominant effect of a single IO infusion of HSD, in agreement with our previous observations (29). In addition the nature of the device prevents complications such as sternal perforation or mediastinitis that have been previously associated with sternal IO infusion (1). Thus, extravasation of fluid into the skin over the sternum was not an observed complication in this study. However, the effects of multiple IO infusions with hypertonic infusions remain to be determined.

In conclusion, the present study indicates that IO infusion of HSD results in rapid delivery of dextran to the vasculature and expansion of plasma volume as effectively as IV infusion. In addition, the IO route did not appear to alter early phases of plasma dextran clearance and did not induce significant pathology to the sternum or lung. Thus, IO infusion of HSD may be a viable alternative for resuscitation from hypovolemia under conditions where access to peripheral vessels is compromised.

Acknowledgement

The authors thank Mr. Donald L. Calkins for preparation of the manuscript and the assistance of the Operating Room Staff of the Division of Military Trauma Research.

REFERENCES

1. Rosetti VA, Thompson BM, Miller J, et al: Intraosseous infusion: An alternate route of pediatric intravascular access. *Ann Emerg Med* 1985; 14:885-888.
2. McNamara RM, Spivey WH, Sussman C: Pediatric resuscitation without an intravenous line. *Am J Emerg Med* 1986; 4:31-33.
3. Iserson KV, Criss E: Intraosseous infusions: A usable technique. *Am J Emerg Med* 1986; 4:540-542.
4. Glaeser PW, Losek JD: Emergency intraosseous infusions in children. *Am J Emerg Med* 1986; 4:34-36.
5. Fiser DH: Intraosseous infusion. *New Engl J Med* 1990; 322:1579-1581.
6. Hodge III D: Intraosseous infusions: A review. *Pediatr Emerg Care* 1985; 1:215-218.
7. Walsh-Kelley CM, Berens RJ, Glaeser PW, et al: Intraosseous infusion of phenytoin. *Am J Emerg Med* 1986; 4:523-524.
8. Schoffstall JM, Spivey WH, Davidheiser S, et al: Intraosseous crystalloid and blood infusion in a swine model. *J Trauma* 1989; 29:384-387.
9. Spivey WH, Lathers CM, Malone DR, et al: Comparison of intraosseous, central and peripheral routes of sodium bicarbonate administration during CPR in pigs. *Ann Emerg Med* 1985; 14:1135-1140.
10. Morris RE, Schonfeld N, Haftel AJ: Treatment of hemorrhagic shock with intraosseous administration of crystalloid fluid in the rabbit model. *Ann Emerg Med* 1987; 16:1321-1324.
11. Tocantins LM, O'Neill JF: Infusions of blood and other fluids into the general circulation via the bone marrow. *Surg Gyn Obstet* 1941; 73:281-287.

12. Tocantins LM, O'Neill JF, Price AH: Infusions of blood and other fluids via the bone marrow in traumatic shock and other forms of peripheral circulatory failure. *Ann Surg* 1941; 114:1085-1092.
13. Bay BK, Henderson JM, Blaisdell FW, et al: A device for rapid vascular access to the sternal marrow spaces for delivery of resuscitation fluids. *Circ Shock* 1989;27:344-345.
14. Watson JC, Pascual JMS, Runyon DE, et al: Intraosseous resuscitation from hemorrhage: Restoration of cardiac output using normal saline (NS) and 7.5% hypertonic saline 6% dextran (HSD). *Circ Shock* 1990; 31:69.
15. Halvorsen L, Bay BK, Perron PR, et al: Evaluation of an intraosseous infusion device for the resuscitation of hypovolemic shock. *J Trauma* 1990;30:652-658.
16. Roe JH: The determination of dextran in blood and urine with anthrone reagent. *J Biol Chem* 1954;208:889-896.
17. Gibson JG Jr, Evans WA Jr: Clinical Studies of blood volume; Clinical application of a method employing the azo dye "Evans Blue" and the spectrophotometer. *J Clinic Invest* 1937;16:301-316.
18. Dubick MA, Summary JJ, Ryan BA, et al: Dextran concentrations in plasma and urine following administration of 6% Dextran-70/7.5% NaCl to hemorrhaged and euvoletic conscious swine. *Circ Shock* 1989;29:301-310.
19. Netter J, Wasserman W: *Applied Linear Statistical Models*. Homewood, IL, Richard D. Irwin, Inc, 1974.
20. Tocantins LM, O'Neill JF, Jones HW. Infusion of blood and other fluids via the bone marrow. *J Am Med Assoc* 1941;117:1229-1234.
21. Meyer LM, Perlmutter M. The absorption rate from the bone marrow. *J Am Med Assoc* 1943;205:187-190.

22. Tocantins LM. Rapid absorption of substances injected into the bone marrow. Proc Soc Exp Biol Med. 1940;45:292-296.
23. Hodge D III, Delgado-Paredes C, Fleisher G: Intraosseous infusion flow rates in hypovolemic "pediatric" dogs. Ann Emerg Med 1987;16:305-307.
24. Maningas PA, Bellamy RF. Hypertonic sodium chloride solutions for the prehospital management of traumatic hemorrhagic shock: A possible improvement in the standard of care? Ann Emerg Med 1986;15:1411-1414.
25. Wade CE, Hannon JP, Bossone CA, et al: Resuscitation of conscious pigs following hemorrhage: Comparative efficacy of small volume resuscitation. Circ Shock 1989;29:193-204.
26. Chavez-Negrete A, Majluf S, Arguero R: Effectiveness of an intraosseous infusion of hypertonic saline dextran during the control of hypotension associated with brain death of organ donors. Eur Surg Res 1990;22:313.
27. Chavez-Negrete A, Majluf S, Frat MA, et al: Treatment of hemorrhagic shock with intraosseous or intravenous infusion of hypertonic saline/dextran solution. Eur Surg Res 1990;22:315.
28. Tocantins LM, O'Neill JF. Complications of intraosseous therapy. Ann Surg 1945;122:266-277.
29. Kramer GC, Mertens SC, Halvorsen L, et al: Intraosseous infusion of hypertonic saline dextran: Effects on pulmonary function and the histology of the bone marrow. Circ Shock 1989; 27:348.

Table 1
 IO vs. IV Infusion of HSD on Plasma Protein Concentrations and Hematocrit¹

	Protein (g/dl)		Hematocrit (%)	
	<u>IO</u>	<u>IV</u>	<u>IO</u>	<u>IV</u>
BL				
0	5.62±0.23	5.34±0.22	26.0±1.3	27.4±1.2
1	4.65±0.14	4.59±0.13	20.0±1.0	21.9±1.8
2	4.58±0.19	4.57±0.15	20.8±1.2	21.4±0.8
4	4.63±0.19	4.68±0.16	21.6±1.3	20.5±0.7
6	4.80±0.20	4.53±0.17	21.8±1.3	22.5±0.9
10	4.75±0.22	4.48±0.23	22.1±1.5	22.8±1.2
15	4.92±0.17	4.62±0.15	22.0±1.5	23.0±1.1
30	4.56±0.26	4.73±0.12	21.5±1.7	22.7±1.2
45	4.85±0.25	4.65±0.14	22.7±1.6	23.3±1.2
60	4.75±0.27	4.60±0.14	22.8±1.7	23.8±1.1
90	4.87±0.25	4.58±0.08	22.9±1.9	24.4±1.3
120	4.95±0.26	4.67±0.14	23.6±2.1	26.0±1.3
	5.00±0.24	4.64±0.12	26.6±1.6	25.4±1.6

¹Data expressed as mean ±S.E. n=6 at each time point except n=5 at 15 min

²BL-Baseline values

³0 min denotes end of infusion period.

Table 2

IO vs. IV Infusion of HSD on Plasma Electrolytes¹

Time (min)	Na (mEq/l)		K (mEq/l)		Cl (mEq/l)	
	IO	IV	IO	IV	IO	IV
BL ²	144±1	142±2	4.7±0.1	4.4±0.1	106±2	105±1
O ₃	169±3	165±6	4.3±0.3	4.0±0.2	136±6	129±6
1	161±3	158±2	4.2±0.3	3.8±0.1	126±2	123±1
2	160±2	158±2	4.2±0.3	3.8±0.1	123±3	121±1
4	158±2	155±2	4.0±0.2	3.6±0.1	122±4	120±2
6	156±2	154±2	4.1±0.2	3.7±0.1	120±4	118±2
10	154±1	152±1	4.2±0.2	3.7±0.1	118±2	117±2
15	154±1	153±1	4.1±0.2	3.8±0.2	119±2	119±1
30	152±1	151±1	4.4±0.2	4.0±0.1	117±4	114±2
45	152±1	150±1	4.4±0.2	4.0±0.1	117±3	113±1
60	152±1	150±2	4.5±0.2	4.2±0.1	116±3	113±2
90	151±1	149±2	4.5±0.1	4.2±0.1	115±2	111±2
120	151±1	149±1	4.4±0.2	4.3±0.1	115±3	110±2

¹Data expressed as mean ±S.E.; n=6 at each time point except n=3 at 15 min²BL Baseline values³O min denotes end of infusion period.

For Na, all times after infusion significantly different (p<0.05) from BL.

Table 3

Histological Observations Following IO or IV Infusion of HSD in Pigs

<u>Treatment</u>	<u>Animal#</u>	<u>Lung</u> <u>Inflammation</u> ¹	<u>Thrombosis or Embolism</u> ²	<u>Sternum</u> <u>Hemorrhage</u> ¹	<u>Washout</u> ^{1,3}
Intraosseous	1	0	1/8	2	0
	2	1	2/6	3	0
	3	1	0	2	1
	4	1	0	0	0
	5	0	0	1	0
	6	0	0	2	0
Intravenous	1	0	0	3	0
	2	0	1/6	0	0
	3	1	2/5	0	1
	4	1	0	2	0
	5	1	1/6	0	0
	6	0	2/6	0	0

¹Data expressed as severity grade where 0= no lesion; 1= minimal; 2= mild; 3= moderate; 4= marked; 5= severe

²Data represent number of sections with thrombi or emboli out of the number of sections examined. At least 5 sections were examined for each animal.

³Washout denotes washout of cellular marrow elements.




Fig 1. Effects of IO and IV infusion of HSD on A) mean arterial pressure, B) cardiac output, C) plasma volume expansion. Data represent mean \pm S.E. of 6 matched experiments.




Fig 2. Plasma dextran concentrations following IO or IV infusion of 4 ml/kg HSD. Data represent mean \pm S.E. of 6 matched experiments.

OFFICIAL DISTRIBUTION LIST

Commander

US Army Medical Research
& Development Command
ATTN: SGRD-RMS/Mrs. Madigan
Fort Detrick, MD 21701-5012

Defense Technical Information Center
ATTN: DTIC/DDAB (2 copies)
Cameron Station
Alexandria, VA 22304-6145

Office of Under Secretary of Defense
Research and Engineering
ATTN: R&AT (E&LS), Room 3D129
The Pentagon
Washington, DC 20301-3080

DASG-AAFJML
Army/Air Force Joint Medical Library
Offices of the Surgeons General
5109 Leesburg Pike, Room 670
Falls Church, VA 22041-3258

HQ DA (DASG-ZXA)
WASH DC 20310-2300

Commandant
Academy of Health Sciences
US Army
ATTN: HSHA-CDM
Fort Sam Houston, TX 78234-6100

Uniformed Services University of
Health Sciences
Office of Grants Management
4301 Jones Bridge Road
Bethesda, MD 20814-4799

US Army Research Office
ATTN: Chemical and Biological
Sciences Division
PO Box 12211
Research Triangle Park, NC 27709-2211

Director
ATTN: SGRD-UWZ-L
Walter Reed Army Institute of Research
Washington, DC. 20307-5100

Commander
US Army Medical Research Institute
of Infectious Diseases
ATTN: SGRD-ULZ-A
Fort Detrick, MD 21701-5011

Commander
US Army Medical Bioengineering Research
and Development Laboratory
ATTN: SGRD-UBG-M
Fort Detrick, Bldg 568
Frederick, MD 21701-5010

Commander

US Army Medical Bioengineering
Research & Development Laboratory
ATTN: Library
Fort Detrick, Bldg 568
Frederick, MD 21701-5010

Commander
US Army Research Institute
of Environmental Medicine
ATTN: SGRD-UE-RSA
Kansas Street
Natick, MA 01760-5007

Commander
US Army Research Institute of
Surgical Research
Fort Sam Houston, TX 78234-6200

Commander
US Army Research Institute of
Chemical Defense
ATTN: SGRD-UV-AJ
Aberdeen Proving Ground, MD 21010-5425

Commander
US Army Aeromedical Research
Laboratory
Fort Rucker, AL 36362-5000

AIR FORCE Office of Scientific
Research (NL)
Building 410, Room A217
Bolling Air Force Base, DC 20332-6448

USAF School of Aerospace Medicine
Document Section
USAFSAM/TSKD
Brooks Air Force Base, TX 78235-5301

Head, Biological Sciences Division
OFFICE OF NAVAL RESEARCH
800 North Quincy Street
Arlington, VA 22217-5000

Commander
Naval Medical Command-02
Department of the Navy
Washington, DC 20372-5120